

High quality cassava flour and industrial starches

Solution Holder is **Abass Adebayo** and can be contacted through **a.abass@cgiar.org**

Summary

Fresh cassava roots are very quick to perish and mold because of their high water content, thereby posing a major challenge for farmers to store the food or sell it on markets. African communities growing cassava have always been processing the roots in one way or another to extend the shelf life and eliminate toxic cyanide compounds. The techniques that are traditionally used for producing cassava flour do not provide significant market opportunities to the smallholder cassava farmers. High quality cassava flour (HQCF) is made through a series of steps, but is not fermented like fufu or gari, making it odorless and giving it a white or off-white color. The main advantages of processing fresh roots into HQCF are the reduction in transport costs from farms to factories, and the longer shelf life of the raw food product. HQCF and derived starches are suitable for manufacturing a wide range of foods and goods, and substitutes imported wheat or other sources of starch. Building capacity in remote rural communities for production of HQCF and starches will boost the performance, value addition and competitiveness along the entire cassava value chain, and thus give rise to societal and economic progress.

Technical Description

Manufacturing HQCF is done using fresh roots with high dry matter and starch content, and low levels of bitter tasting cyanides, and needs to take place within a day or two after harvest so the desired quality is obtained. The key steps in producing HQCF are: raw material selection, peeling, washing, grating, pressing, disintegration, sifting, drying, milling, screening, packaging, and storage; though slicing or chipping can be performed instead of grating, pressing, disintegration and sifting. Detoxification of cassava roots in this process happens through grating, dewatering, and drying that unlike fermentation do not result in a bitter taste and odor. Extraction of starch from cassava follow a similar process expect that it requires wet milling of grits or chip and physical separation of fibers. To achieve a high degree of palatability and starch content in the flour product it critical that fresh cassava roots get delivered on time and are processed with appropriate methods, for which capacity needs to be built within farmer communities. HQCF is a gluten-free product, making it ideally suited for manufacturing of non-allergenic foods which a major growth market globally.

Uses

HQCF is suitable for manufacturing a large range of food product, such as soups, bread, glucose syrups and alcohol. The proportion of HQCF that can be used in bakery products amounts from 10% to 35%, and the heavy cake structure bread is preferred in rural areas. Through the use of enzymes from plant seedlings it is possible to convert of HQCF into sugar syrups which are similar to common dextrose products and meet different end-user requirements. In further steps, the sugar syrup derived from cassava can be made into alcohol for potable and industrial use. When blending HQCF with salt and soda it can be used as adhesive for paper board and plywood glues, and extracted starches as filler for cosmetic, pharmaceutical and regular products, or as textile stiffener; completely replacing imported starch materials derived from wheat and other crops.

Composition

Nutritional and chemical properties of HQCF or derived starches are depending on the used cassava variety, freshness of roots and processing steps; all of which have to be harmonized with the customer requirements. The typical quality standards of HQCF are: moisture content 13%, fiber 2%, ash 3%, fine flour (<0.6 mm) 90%, hydrocyanic acid 10 mg/kg, sulfated ash 0.5%, starch content 65 - 70%, protein content 1.3 - 2.0%, crude fat content 0.3 - 0.5%, and total acidity 1%. Pro-vitamin A or beta-carotenoids levels measure 0.06 µg/g for HQCF that is made from white fleshed cassava, whereas this goes up to 10.7 µg/g when biofortified yellow fleshed cassava is used to prepare HQCF. The moisture content of flour determines its storage stability, as the lower the flour moisture, the higher the storage stability. The cyanide content for HQCF is very low (2.04 to 48.13 ppm) and within safe level for human consumption. Foaming and emulsion capacities of HQCF is low but its bulk densities, water and oil absorption capacities as well as degree of gelatinization are high.

Means of application

Manufacturing of HQCF can be done with simple cottage style equipment that are manually operated, similar to that already used for gari processing. Large automated systems are available for industrial scale processing of cassava roots into flour. The minimal infrastructure required for manufacturing of HQCF include: a grater for size reduction, press for dewatering, solar or flash dryer for drying, pin mill for comminution and stitching machine for packaging. For production of starch there is need to remove cellulose fibers that were released in the wet milling step, which can be done with a jet extractor or perforated-basket centrifuge. Reliable supply of low-cost electricity from the national grid, diesel generators or photovoltaic panels has to be in place for warranting uninterrupted and cost-effective manufacturing of HQCF and starches.

| | |
|----------------------|-------------------------|
| Agroecologies | All Agroecologies. |
| Regions | Africa South of Sahara. |

| | |
|---------------------------------|--|
| Developed in Countries | Nigeria. |
| Available in | Nigeria. |
| Solution Forms | Other. |
| Solution Applications | Post-harvest handling, Value addition, Agri-Food Processing. |
| Agricultural Commodities | Cassava. |
| Target Beneficiaries | Small-scale farmers, Commercial farmers. |

Commercialization

Commercialization Category

Commercially available

Startup Requirements

1) Raise awareness with cassava farmers, agri-food companies and goods manufacturers in the country about the benefits of HQCF and starch, 2) Identify profitable, durable and equitable integration of HQCF and starch in local markets, 3) Organize reliable supply of cassava roots with high dry matter and starch content close to the processing plant, 4) Set up processing methods and simple energy-efficient and labor-saving equipment such as graters, and dewatering machines, and 5) Train machine operators and workers on quality adherence throughout the manufacturing process

Production Costs

The cost of cassava fresh roots at the factory gate, including transportation, is a major determinant of the economic viability of manufacturing HQCF and derived starches. Producing 1 ton of HQCF requires on average 5.5 ton of fresh cassava roots. Approximately 60% of the capital investment that is required to set up a processing plant is going to machinery and equipment, while the rest to construction. A demonstrator carried out in Madagascar showed that processing 288MT of fresh cassava roots into HQCF over a year costed USD 17,238, which corresponds to a production cost of USD 60 per MT fresh root product. Production of starch is more capital intensive and requires higher levels of inputs as compared to HQCF, making larger investments are needed to enter that market.

Customer Segmentation

smallholder farmers, farmers association, farmers cum-processors, industrial flour processors, and food manufacturers.

Potential Profitability

A study in Uganda on production of HQCF found that a net margin of USD 78.9/ton can be achieved when starting from dried chips, which suggests a real potential for enterprise development in remote rural communities. HQCF is currently sold at USD 550-650/ton and represents a potential 25% reduction in raw material costs for bakeries that now use wheat flour; e.g., for a producer of biscuit that processes 200MT flour per month, the substitution of wheat flour by HQCF can lead to an annual saving of USD 130,560 per annum. The higher value of HQCF and starch than traditionally prepared cassava flour makes that farmers will earn additional incomes, and can venture into local manufacturing.


Licensing Requirements


Producers of HQCF and starch have to comply with food safety regulations. Most of the simple cottage style machinery and equipment can be fabricated free of license, while industrial systems fall under intellectual property protection rules.

Innovation as Public Good

The International Institute of Tropical Agriculture is responsible for the public development of high quality cassava flour technologies that are released to countries in Sub-Saharan Africa.

Solution Images





Institutions



Accompanying Solutions

Golden cassava varieties, Cassava varieties with high dry matter and starch content,
Mobile cassava processing plant