

## **In-Vitro Tissue Culture Propagation**

Solution Holder is **John Derera** and can be contacted through **[j.derera@cgiar.org](mailto:j.derera@cgiar.org)**

### **Summary**

Banana and plantain is propagated in the laboratory through tissue culture (TC). In vitro micro-propagation eliminates all pests and diseases except for viruses. TC consists of five important steps: initiation, multiplication, shooting and rooting, primary hardening in green houses and secondary hardening in shade houses. TC plants have the benefits of uniformity and fast propagation of large numbers of plantlets. These advantages enable marketing and more rapid recovery from broad-scale damage such as disease outbreak and extreme weather.

### **Technical Description**

In-vitro micro-propagation relies upon a liquid, semi-solid, or solid growth medium such as broth or agar multiplied under sterile growing conditions. The most common tissue culture techniques is meristem culture but other techniques such as callus culture, somatic embryogenesis, cell suspension, and protoplast culture are possible. TC plantlets tend to be less robust than suckers and require greater care due to their initial sterile production devoid of beneficial microorganisms. They are nevertheless regularly planted into fields with high pest and disease burdens and abiotic constraints. Inoculation of TC plantlets with beneficial microorganisms such as *Fusarium oxysporum* and *Trichoderma* spp. helps to reintroduce immunity. Micro-propagation through tissue culture techniques is an efficient method of producing large quantities and good quality plantlets but requires high capital investment and skills. Strict adherence to aseptic standards and micro-climatic conditions and care during the hardening is important. Careful examination of the stock is required to avoid culturing the infected plants. Farmers should be aware that planting TC culture plantlets from a single banana or plantain species causes low genetic diversity and variability that poses a risk if disease or pest outbreak occurs.

### **Uses**

Tissue-cultured banana seedlings are not always conveniently available. Larger-scale banana farmers may wish to establish their own banana tissue-culture facility to ensure availability of disease-free seedlings. To make TC plantlets readily accessible to farmers in remote areas, there is need for linkages with macro-propagation nurseries.

### **Composition**

Water purification systems for producing double distilled water are needed to prepare the stock solutions and media, autoclave for sterilizing tools and media, pH meter and refrigerator to store the stock solutions and hormones. Laminar airflow cabinets to perform culturing and subculturing plants, these can be high-end models for specialized laboratories with large output capacity but may also be fabricated with plastic boxes and a small air filter. Other materials required include a pH meter, hot plate/stirrer, glasswares and beakers, forceps, bunsen burner, wash bottles, brushes, culture vessels and tubes, Erlenmeyer flasks, graduated cylinders, glass pipettes, scalpel handles, and blades, spatulas, stir bars, roll tape, gloves, parafilm, lab markers, paper towels, detergents, isopropyl alcohol, chlorine bleach, and brushes. Growth media for the preculture are made with Murashige & Skoog basal medium supplemented with 5.0 mg/l BAP, and subculture media are half of those concentrations. An air-conditioned room fitted with two 40-watt fluorescent tubes and a 16-hour photoperiod.

### Means of application

The first step is to collect suckers 40–100 cm tall from strong mother plants that are free of disease symptoms, making sure to separate the sucker without cracking the corm. It is advised to collect at least two suckers from each plant source, one for micropropagation and the other for a nursery farm for future propagation. Approximately 10 cm of inner tissue containing the meristematic corm is excised from the stem by removing the leaf sheaths in a clean laboratory and rinsed with water to wash of soil. The second part of the process happens in a laminary flow hood using autoclaved scalpels, forceps, cutting plates and solutions. The entire corm is disinfected by submerging it in series of disinfectant solutions. Next, the surface-sterilized corm is trimmed until it measures 1 × 1 cm, with the corm tissue as thin as possible. On a fresh cutting plate, the corm is cut longitudinally into propagules of 0.5 cm. Propagules are placed in tubes with growth media and closed with an open cap. Over one month the propagules are kept in the room during which bulging and browning of the corm, greening of leaf tissue and shoot emergence is monitored. Contaminated or off-growth precultures must be discarded. When the propagules are 2 cm tall, they are transferred from tubes into jars with growth media to proliferate more shoots. Larger propagules can be cut in two. These subcultures must be kept in the growth chamber for 3-4 weeks to obtain desired numbers of shoots.

<b>Agroecologies</b>	Highlands, Humid forest, Moist savanna.
<b>Regions</b>	Africa South of Sahara.
<b>Developed in Countries</b>	Zambia, Uganda, Tanzania, Somalia, Rwanda, Nigeria, Kenya, Ivory Coast, Ghana, Ethiopia, Democratic Republic of the Congo, Cameroon, Burundi.

<b>Available in</b>	Zambia, Uganda, Tanzania, Somalia, Rwanda, Nigeria, Kenya, Ivory Coast, Ghana, Ethiopia, Democratic Republic of the Congo, Cameroon, Burundi.
<b>Solution Forms</b>	Input Supply.
<b>Solution Applications</b>	Vegetative propagation.
<b>Agricultural Commodities</b>	Banana/Plantain.
<b>Target Beneficiaries</b>	Commercial farmers, Agro-dealers.

## Commercialization

### Commercialization Category

Commercially available

### Startup Requirements

Steps needed for starting a TC production plant involve: 1) Business planning and proper market analysis, 2) Obtaining a loan from a bank or other financing institution to acquire equipment, 3) Training of operating staff on handling and quality control procedures, and 4) Awareness raising of nearby farmers about planting and macro-propagation of TC plantlets.

### Production Costs

The culturing procedure of the banana and plantain differs between species. So, to establish a new TC product offer, several experiments and trial and error methods must be performed. High cost of plantlet production through micropropagation technique is a major concern limiting its wide application, despite its obvious advantages. Improving process efficiency and better utilization of resources can lower the cost of production without compromising the quality of the micropropagules and plants.

### Customer Segmentation

In Africa, a number of commercial tissue culture laboratory enterprises are operating that supply farmers with clean high quality material for planting and macro-propagation.

### Potential Profitability

The efficiency of in-vitro propagation systems is mainly influence by the rate of multiplication which is dependent on genotype as well as operating procedures. Plantlets

are sold for about US \$1.3 to \$1.5 by large commercial retailers. A nursery business can produce more than 3,000 TC plantlets per cycle and earn a gross revenue of US \$1,600, achieving a return on investment of 1.4. The benefit of TC plantlets is that stands are uniform and follow the same growth timeline so they can be harvested at the same time which simplifies plantation management and reduces overall costs. Where TC material is used, farmers shorten production cycles allowing earlier harvest and allowing more than 90% of plants bear bunches.


### **Licensing Requirements**

In some cases, micro-propagation techniques are licensed but mostly they are protected by trade secrets. Once purchased, producers are generally free to multiply their stock using macro-propagation techniques.

### **Innovation as Public Good**

In-vitro propagation techniques are disseminated as Public Good.

# Solution Images



*Steps of in-vitro tissue culture micro propagation: a) Removal of sheaths, b) Separated corm, c) Desinfection and*

## Institutions



## Accompanying Solutions

[Improved Varieties of Plantain for Tropical Lowlands](#)

[Improved Varieties of Banana for the African Highlands](#)

[Propagation of Disease-Cleaned Suckers](#)